
Climate Change in the Mediterranean. The Jucar River Basin Case

Miguel Ángel Pérez-Martín

mperezm@hma.upv.es

<http://www.upv.es>

Malta 23 March 2017

1




I) Observed data

2




Temperature and sea level anomaly

Mean global Temperature ocean-land
 Reference Period 1960-1990
 2015 +0,87°C Record
 2016 +0.99°C Record




Earth observatory NASA
<http://climate.nasa.gov/>
<http://earthobservatory.nasa.gov/>



Mean Temperature Anomaly 2000-2009

Data from ground stations 1870-2000 (130 years)
 Increase of 20 cm, from 1870
 1.5 cm per decade CSIRO

Data from satellite 6 cm from 2000
 3.5 cm per decade (NASA)





CO2 and Methane levels in the atmosphere

CO2
 Half-life
 100 years

Year 1960 (320 ppm).
 Year 2017 (>400 ppm)



<http://www.esrl.noaa.gov/gmd/ccgg/trends/>



Methane
 Half-life
 12 years

Year 1980 (1650 ppb)
 Year 2017 (1850 ppb)


http://www.esrl.noaa.gov/gmd/ccgg/trends_ch4/



CO2 Reconstruction last 2.000 years (Roman Age)

Main Greenhouse Gases:
 Carbon Dioxide CO₂
 Methane CH₄
 Nitrous Oxide N₂O

Third Report
 2001




CO2 Preindustrial (270 ppm). Year 1960 (320 ppm). Year 2017 (>400 ppm.)

Paleoclimate: 600.000 years Reconstruction (Discovery of Fire)

Quaternary
 2.6 Millions of
 Years
 Discovery of Fire
 by Humans
 400.000 years


600.000 years 130.000 years



Deuterium δD
 Hydrogen Isotope


- Temperature an CO2 well correlated.
- Calide Periods: 240-300 ppm. Glacial Periods: 180-200 ppm

Current Day




II) Scenarios and models

Name	Radiative Forcing ¹	Concentration ²	Pathway shape
RCP8.5	>8.5 W/m ² in 2100	> ~1370 CO ₂ -eq in 2100	Rising
RCP6	~6 W/m ² at stabilization after 2100	~850 CO ₂ -eq (at stabilization after 2100)	Stabilization without overshoot
RCP4.5	~4.5 W/m ² at stabilization after 2100	~650 CO ₂ -eq (at stabilization after 2100)	Stabilization without overshoot
RCP3-PD ³	peak at ~3W/m ² before 2100 and then decline	peak at ~490 CO ₂ -eq before 2100 and then decline	Peak and decline





Radiative Forcing of the Representative Concentration Pathways. van Vuuren et al (2011)
The Representative Concentration Pathways: An Overview. Climatic Change, 109 (1-2), 5-31.




Temperature and sea level results


Temperature:
RCP2.6 1°C
RCP4.5-6 2°C
RCP8.5 4°C




Temperature scaled by global T (°C per °C)



Precipitation scaled by global T (% per °C)




Sea level increase
40 cm RCP2.6
60 cm RCP8.5
Max 1.0 m




EC 2007. Green Paper from the Commission to the Council, the European Parliament, The European economic and social Committee and the Committee of the Regions.
Adapting to climate change in Europe – options for EU action

Cambios en la temperatura



Cambios en la precipitación





Previsiones de variación para 2071-2100 referido al periodo 1960-1990
Basado en: IPCC SRES escenario A2. Proyecto PRUDENCE


North Europe:

2030-2050: ΔT 2°C ΔP +5+10%
2080-2100: ΔT 4°C ΔP +10+15%

Scenario A1B




Projected mean annual temperature (°C) and precipitation (%) change relative to 1961-1990




Mediterranean Area:

2030-2050: ΔT 2°C ΔP -5-10%
2080-2100: ΔT 4°C ΔP -15-20%



European 0
Precipitation Line
Scenario RCP 8.5
Year 2100



White Zone: -5% a 5%
Change in Precipitation



III) Impacts in the Júcar River Basin


- Reduction in the natural water resources (average conditions)
- Increase of number and intensity of Droughts (extreme conditions)
- Increase in the Risk of Wildfire in the river basin (extreme conditions)
- Water needs for crops (average conditions)



Water Balance Model Patrical

Monthly Distributed Water Balance Model
Surface+ Groundwater+
River-aquifer interaction






- **MPI-ESM-MR.**- Max-Planck-Institut-Earth System Model running on médium resolution grid
- **HadGEM2-ES.**- Hadley Global Environment Model 2 - Earth System


Climate Change Impact on Water Resources and Droughts of AR5 scenarios in the Jucar River, Spain Miguel A. Pérez-Martín, A. Batán, P. del-Amo , S. Moll. ICDrought 2015

Temperature and Precipitation Anomalies refer to 980-2005 period




- Based on: **AR4 IPCC, 2007**.
- Dynamic downscaling: **ECHAM5, T63 (1.9°) L31**
- Scenario: **A1B**
- Regional model: **RegCM3**
- Control period: 1990 - 2000
- **Future period: 2010 - 2040**

Reduction natural resources:
 River flows 21%
 In the headers (Alarcón 26%, Contreras 28%)
 Rainfall recharge into aquifers: 19%




Doctoral Thesis: "Caracterización de los futuros escenarios climáticos en la Comunidad Valenciana: propuestas de mejora para la evaluación de la oferta y demanda de recursos hídricos".
 Chirivella Vicente, 2010. Directores: José Capilla y Miguel Ángel Pérez

Temperature





Variaciones en las aportaciones (2070-2100)



SRES A2. 2070-2100

Precipitation



Escenarios hidrológicos futuros:

Escenarios climáticos 2070-2100:

$\Delta T \approx +3^\circ\text{C}, +4^\circ\text{C}$

$\Delta P \approx -15\%, -25\%$

Efectos escenarios


hidrológicos 2070-2100:

$\Delta Q \approx -30\%, -40\%$

Doctoral Thesis: "Efectos del cambio climático en los sistemas complejos de Recursos Hídricos. Aplicación a la cuenca del Júcar". Hernández Barrios, 2007. Directores Joaquín Andreu y Miguel Ángel Pérez

Mediterranean one of the most vulnerable areas


Variaciones porcentuales en las escorrentías anuales periodo 2090-2099, respecto periodo 1980-1999. Escenario A1B



Spain reductions around 40% 2090-2099


Zonas ralladas: más de un 90% en la concordancia de los modelos en el signo del cambio "Climate Change and Water", IPCC Technical Paper VI. IPCC", june 2008 19

Increase of number and intensity of Droughts



Palmer Index

Drought Definition
Moderate (-2)
Severe (-3)
Extreme (-4)




Probability of Drought with Socio-Economical Consequences (Severe):

- Double in the short-term (2010-2030)
- 6-9 times in the long-term (2080-2100)

Climate Change Impact on Water Resources and Droughts of AR5 scenarios in the Jucar River, Spain. Pérez-Martín, Batán, del-Amo and Moll. ICDrought 2015

Risk of Wildfires and Potential number



Canadian Fire-Weather Index (FWI) (Van Wagner, 1974)



- Increase in the number of potential Wildfires per year:
- 40% in the short-term
 - Doubling in the long-term

Risk level	FWI	KBDI	Potential number of Wildfires per season
Low	0 – 4	0 – 40	0
Medium	4 – 18	40 – 180	20 – 60
High	18 – 28	180 – 280	60 – 150
Very High	28 – 39	280 – 390	150 – 300
Extreme	>39	>390	+300

Pérez-Martín, M.A., Vinces, F., del-Amo, P., Batán, A. 2015. *Climate Change and the Increase of Wildfire Risk in the Upper Júcar River Basin, Spain. Drought: Research and Science-Policy Interfacing - Proceedings of the International Conference pp. 91-96*

Water needs for Crop Irrigation

Cropwat software
Penman-Monteith method






Citric: current: 4.000-4.500 m³/ha/year
Future: 6.000-6.500 m³/ha/year
increase: 40%-50%


Mean water needs increase for all irrigation crops 2100: 30% - 40%

Doctoral Thesis: "Efectos del cambio climático en los sistemas complejos de Recursos Hídricos. Aplicación a la cuenca del Júcar". Hernández Barrios, 2007. Directores Joaquín Andreu y Miguel Ángel Pérez

Productivity Increase in Cold Regions ($T < 15^{\circ}\text{C}$)
 Productivity Reduction in Warm Regions ($T > 15^{\circ}\text{C}$)
 Spain: $T_{\text{mean}} 15^{\circ}\text{C}$. North 14°C , Mediterranean $15\text{-}18^{\circ}\text{C}$



Based on:
 Laboral supply
 Labour Performance
 Agricultural Yield



Marshall Burke, Solomon M. Hsiang & Edward Miguel 2015. Global non-linear effect of temperature on economic production. Nature

IV) Conclusions

Conclusions about future scenarios

- Changes in climate in the Mediterranean area, Jucar Basin:
 - Temperature Increase: **1°C next decades** (2010-2030) and **3 - 5 °C at the end of the century.**
 - Rainfall Reductions: **10% next decades** (2010-2030) and 20% at the end of the century.
- Impacts:
 - Natural Water Resources will be reduced, **20% next decades** (2010-2030) and more than **40%** (aprox. 50%) (2070-2100).
 - **Droughts will be increased** in number and severity: **doubling** in short-term and 6 or more times in long-term
 - The **Number of Wildfires per year** will be **increased 40%** next decades (2010-2030) and doubling at the end of the century.
 - **Irrigation water requirements will increase around 40%.**
- **Mediterranean area** is one of the more **vulnerable** areas to the Climate Change in relation with the Water Resources

Conclusions about future scenarios

- Current CO₂ level 400 ppm future levels from 600-1200 ppm. Return to 60-100 Millions of Years
- Current carbon emissions is curving but is far of the 2° increase pathway. (Curving US gas, and Chine carbon reduction aerosols)
- **Mediterranean. Jucar River Basin:**
 - **Natural water resources reduction and irrigation water demand increase** (with the same crops) will contribute to a **more stressed River Basin.**
 - **Gradual reduction of the Natural Resources.** It is necessary to learn how to preserve the environment and produce the same with the half of water. Increase the efficiency in the water use.
 - **Adaptive measures are required.** Also, **permanent monitoring** of the climate change evolutions is required to reduce the vulnerability of the system.

Thank you for
your attention

Turia river



Albufera wetland



Júcar river